

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>5</sup> :</b> C07K 13/00, A61K 37/02 C12P 21/02, C12N 15/24 C07H 15/12	<b>A1</b>	<b>(11) International Publication Number:</b> WO 91/05798 <b>(43) International Publication Date:</b> 2 May 1991 (02.05.91)
<b>(21) International Application Number:</b> PCT/US90/05522 <b>(22) International Filing Date:</b> 27 September 1990 (27.09.90) <b>(30) Priority data:</b> 420,038 10 October 1989 (10.10.89) US <b>(71) Applicant:</b> AMGEN INC. [US/US]; 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US). <b>(72) Inventors:</b> BOONE, Thomas, C. ; 3913 Elkwood, Newbury Park, CA 91320 (US). MILLER, Allan, L. ; 2111 Balmain Way, Glendale, CA 91206 (US). <b>(74) Agent:</b> ODRE, Steven, M.; Amgen Inc., 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US).		<b>(81) Designated States:</b> AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent)*, DK (European patent), ES (European patent), FR (European patent), GB (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), SE (European patent).  Published With international search report.
<b>(54) Title: COMPOSITIONS AND METHODS FOR TREATING OR PREVENTING INFECTIONS IN CANINE AND FELINE ANIMALS</b>  <b>(57) Abstract</b>  Compositions and methods for treating or preventing infections in canine or feline animals which comprises administering an effective amount of granulocyte colony stimulating factor (G-CSF), are disclosed. The G-CSF may be naturally derived, or alternatively, the G-CSF and genetically engineered variance of G-CSF may be the expression products of genetically engineered prokaryotic or eukaryotic host cells.		

Ala Pro Leu Gly Pro Thr Gly Pro Leu Pro Gln Ser Phe Leu Leu  
gcc ccc ctg ggc cct acc ggc ccc ctg ccc cag ago ttc ctg ctc

Lys Cys Leu Glu Gln Met Arg Lys Val Gln Ala Asp Gly Thr Ala  
aag tgc cta gag caa atg agg aag gtc cag gct gat ggc acg gcg

Leu Gln Glu Thr Leu Cys Ala Thr His Gln Leu Cys His Pro Glu  
ctg cag gag acg ctg tgt gcc acc cac cag ctg tgc cat cct gag

Glu Leu Val Leu Leu Gly His Ala Leu Gly Ile Pro Gln Pro Pro  
gag ttg gtg ctg ctc ggg cac gct ctg ggc atc ccc cag cct ccc

Leu Ser Ser Cys Ser Ser Gln Ala Leu Gln Leu Met Gly Cys Leu  
ctg agc agc tgc tcc agc cag gcc ctg cag ctg atg ggc tgc ctg

Arg Gln Leu His Ser Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln  
cgt caa ctc cac agc ggc ctc ttc ctc tac cag ggc ctc ctg cag

Ala Leu Ala Gly Ile Ser Pro Glu Leu Ala Pro Thr Leu Asp Thr  
gcc ctg gca ggg ata tcc ccc gag tta gcg ccc acc ttg gac aca

Leu Gln Leu Asp Thr Thr Asp Phe Ala Ile Asn Ile Trp Gln Gln  
ctg cag ctg gac acc acc gac ttt gcc atc aac atc tgg cag cag

Met Glu Asp Leu Gly Met Ala Pro Ala Val Pro Thr Gln Gly  
atg gaa gat cta gga atg gcc ccc gcc gtg cca cct acc cag ggc

Thr Met Pro Ala Phe Thr Ser Ala Phe Gln Arg Arg Ala Gly Gly  
acc atg cca gcc ttc acc tcy gcc ttc cag cgc cgg gca gga ggt

Val Leu Val Ala Ser Asn Leu Gln Ser Phe Leu Glu Leu Ala Tyr  
gtc ctg gtg gcc tcc aac ctg cag agc ttc ctg gag ctg gca tat

Arg Ala Leu Arg His Phe Ala Lys Pro  
cgc gct ctg cgc cac ttt gcc aac ccc

### DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CI	Côte d'Ivoire	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

- 1 -

COMPOSITIONS AND METHODS FOR TREATING  
OR PREVENTING INFECTIONS IN CANINE AND FELINE ANIMALS

Field Of The Invention

5

The present invention is directed to the use of granulocyte colony stimulating factor (G-CSF) to treat or prevent infections in canine and feline animals. More specifically, the invention is directed to the use of G-CSF having the amino acid sequence of human G-CSF or having the amino acid sequence of canine G-CSF, in treating or preventing infections in canine or feline animals. The source of the G-CSF may be naturally derived or may be derived from genetically engineered prokaryotic or eukaryotic host cells containing recombinant plasmid or viral DNA vectors carrying the human or canine G-CSF gene, or genetically engineered variants of canine G-CSF genes, or synthetic human or canine G-CSF genes. The present invention is also directed to DNA gene segments, biologically functional recombinant plasmids and viral DNA vectors, and prokaryotic and eukaryotic host cells containing such recombinant plasmids and vectors, all of which contain a canine G-CSF gene or a genetically engineered variant of a canine G-CSF gene.

Background Of The Invention

Although antibiotic therapy is now used for animal infections with some success, huge losses persist. The early hopes that antibiotics would allow complete control of the disease have not been realized. None of the antibiotics utilized thus far has been entirely satisfactory. Additionally, it has been found to be very desirable to replace antibiotic treatment with treatment by non-antibiotic chemotherapeutic drug compounds, for the following reasons:

- 2 -

(1) Antibiotics effective in human medicine should not be utilized in veterinary medicine, in order not to build up strain resistance of bacteria appearing in human diseases; and

5 (2) Antibiotics should be reserved for such diseases for which no chemo-therapeutic drug compound would be available, as it has been proved that bacterial strains build up resistance to an antibiotic after extended use of such antibiotic.

10 Despite these several published methods, it remains very important to find cost-effective methods utilizing non-antibiotic compounds which would substantially overcome the drawbacks of antibiotics used thus far and yet would be effective in treating and  
15 preventing infections in canine and feline animals.

Canine parvo virus still infects over one-half million young dogs. Hospitalization and intensive care are required. Mortality occurs in 15-20% of the cases. Severe neutropenia occurs and death is thought  
20 to frequently result from secondary infections and sepsis.

Feline Immunodeficiency Virus (FIV) is believed to infect 500,000-1,000,000 cats per year. This virus causes neutropenia in approximately 30% of  
25 the cats which renders them susceptible to infections. Feline Leukemia Virus (FeLV) also causes neutropenia in cats.

#### Granulocyte Colony Stimulating Factor

30 Granulocyte colony stimulating factor (G-CSF) is one of several glycoprotein growth factors known as colony stimulating factors (CSFs) because they support the proliferation of haemopoietic progenitor cells. G-CSF stimulates the proliferation of specific bone  
35 marrow precursor cells and their differentiation into granulocytes. It is distinguished from other CSFs by

- 3 -

its ability to both stimulate neutrophilic granulocyte colony formation in semi-solid agar and to induce terminal differentiation of murine myelomonocytic leukemic cells in vitro. The cDNA cloning and expression of recombinant human G-CSF has been described, and it has been confirmed that the recombinant G-CSF exhibits most, if not all, of the biological properties of the native molecule (Souza, L. et al. Science 232, 61-65 (1986)). Sequence analysis of the cDNA and genomic DNA clones has allowed the deduction of the amino acid sequence and reveals that the protein is 204 amino acids long with a signal sequence of 30 amino acids. The mature protein is 174 amino acids long and possesses no potential N-linked glycosylation sites but several possible sites for O-linked glycosylation.

The cloning and expression of cDNA encoding human G-CSF has been described by two groups (Nagata, S. et al., Nature 319, 415-418 (1986); Souza, L. M. et al., Science 232, 61-65 (1986)). The first report of a G-CSF cDNA clone suggested that the mature protein was 177 amino acids in length. The authors reported that they had also identified a cDNA clone for G-CSF that coded for a protein that lacked a stretch of three amino acids. This shorter form of G-CSF cDNA expresses the expected G-CSF activity. The second report describes a cDNA sequence identical to this short form and makes no mention of other variants. Since these authors confirmed that the short cDNA expresses G-CSF with the expected profile of biological activity, it is probable that this is the important form of G-CSF and that the longer form is either a minor splicing variant or the result of a cloning artifact.

Matsumoto et al., in Infection and Immunity, Vol. 55, No. 11, p. 2715 (1987) discuss the protective effect of human G-CSF on microbial infection in neutropenic mice.

- 4 -

The following patent publications relate to G-CSF: WO-A-8703689, assigned to Kirin/Amgen describes hybridomas producing monoclonal antibodies specific for G-CSF and their use in the purification of G-CSF; WO-A-  
5 8702060, assigned to Biogen, discloses human G-CSF like polypeptides and methods of producing them; U.S. Patent 4,810,643 assigned to Amgen, discloses human G-CSF like polypeptides, sequences encoding them and methods of their production; and WO-A-8604605 and WO-A-8604506,  
10 both assigned to Chugai Seiyaku Kabushiki Kaisha, disclose a gene encoding human G-CSF and infection inhibitors containing human G-CSF.

The use of recombinant G-CSF with the same amino acid sequence as human G-CSF, in dogs with cyclic  
15 neutropenia has been associated with the development of neutralizing antibodies to the heterologous G-CSF protein during a thirty day period of administration (see Lothrop et al., Blood 72, 5624-37 (1988)). Subsequent treatment of these same dogs with recombinant  
20 human GM-CSF failed to cause a significant leukocytosis or eliminate cycles of neutropenia. A significant variation in structure may explain the development of neutralizing antibodies when the human sequence products are given to dogs. The development of neutralizing  
25 antibodies in dogs given the human sequence products may limit them to single or short term use.

It is an object of the subject invention to provide an improved method of treating and preventing infections in canine or feline animals.

30 It is a further object of the subject invention to provide a method of treating infections in canine or feline animals without build up of strain resistance of bacteria.

A still further object of the invention is to  
35 provide a purified and isolated polypeptide having part or all of the primary structural conformation and the

- 5 -

biological properties of naturally occurring canine G-CSF, and DNA sequences encoding such G-CSF.

Other objects, features and characteristics of the present invention will become apparent upon  
5 consideration of the following description and the appended claims.

#### Summary Of The Invention

10 The present invention provides DNA sequences, biologically functional recombinant plasmids and viral DNA vectors, and prokaryotic and eukaryotic host cells containing such recombinant plasmids and vectors, all of which contain a canine G-CSF gene or a genetically  
15 engineered variant of a canine G-CSF gene. The invention also provides polypeptides encoded by the canine G-CSF gene or variants thereof. A method for treating or preventing infections in canine or feline animals is also disclosed.

20 Novel DNA sequences of the invention include sequences useful in securing expression in prokaryotic or eukaryotic host cells of polypeptide products having at least a part of the primary structural conformation and the biological properties of naturally occurring  
25 canine granulocyte colony stimulating factor. DNA sequences of the invention are specifically seen to comprise the DNA sequence of the coding region of the mature protein, set forth in Figure 2 or its complimentary strand, allelic variant forms of canine  
30 G-CSF, manufactured DNA sequences encoding canine G-CSF, fragments of canine G-CSF and analogs of canine G-CSF with DNA sequences incorporating codons facilitating translation of messenger RNA in microbial hosts. Such manufactured sequences may readily be constructed  
35 according to the methods of Alton, et al., PCT published application WO 83/04053.

- 6 -

A further embodiment of the invention relates to synthetic genes designed to allow for expression of G-CSF having the canine amino acid sequence in E. coli.

Also comprehended by the invention are

5 pharmaceutical compositions comprising effective amounts of polypeptide products of the invention together with suitable diluents, adjuvants and/or carriers useful in animal therapy.

The subject invention also relates to a method  
10 for treating and preventing infections in canine or feline animals by administering a therapeutically effective treating or preventing amount of granulocyte colony stimulating factor, advantageously G-CSF derived from the gene of a canine animal. In addition, the  
15 invention relates to a method of treating cancer in canine or feline animals by administering a therapeutically effective treating or preventing amount of granulocyte colony stimulating factor as an adjunct to chemotherapy.

20

#### Brief Description of the Drawings

Figure 1 shows the restriction map of canine G-CSF;

25 Figure 2 illustrates the coding region of the mature protein of canine G-CSF;

Figure 3 is the genomic sequence of the human G-CSF;

Figure 4 is the DNA sequence of a canine G-CSF synthetic gene (cG-CSF dna);  
30

Figure 5 illustrates the oligos used to construct the subunits of the canine G-CSF synthetic gene (cG-CSF dna3);

Figures 6A and 6B shows the two subunits of  
35 the canine G-CSF synth tic gene cG-CSF dna3;



- 7 -

Figure 7 shows the homology of canine and human G-CSF.

Figures 8-9 are graphic representations of the results obtained in Example 5 which relates to treatment of dogs with G-CSF having the canine amino acid sequence.

#### Detailed Description Of The Invention

10           A novel method for treating or preventing infections in canine or feline animals has been discovered. Surprisingly it has been found that G-CSF is effective in a method of treating or preventing infections in canine and feline animals.

15           The subject invention also relates to treating cancer in dogs or cats by administration of G-CSF as an adjunct to chemotherapy, advantageously, as an adjunct to the use of myelosuppressive drugs. The general method as it applies to humans is described in Gabrilove  
20 et al., New England Journal of Medicine 318, No. 22 (1988) hereby incorporated by reference. A skilled veterinarian will adjust the method of administering dose etc. as appropriate.

          A variety of infections afflicting canine and  
25 feline animals are treatable by the method of the subject invention. A veterinarian of ordinary skill can readily determine whether an animal exhibits an infection. In one embodiment, the present invention relates to a method of treating or preventing infections  
30 such as Feline Immunodeficiency Virus (FIV) in feline animals comprising administering a composition which comprises an effective amount of G-CSF.

          In another embodiment of the invention, G-CSF is used to treat Feline Leukemia Virus (FeLV).  
35 Additionally G-CSF is used to treat cats with Pan Leukopenia.

- 8 -

In another embodiment dogs infected with Parvo Virus are treated with G-CSF.

The subject invention also relates to the use of G-CSF during bone marrow transplants. G-CSF shortens the time to engraftment (4-7 days vs. 7-10 days in a study with 12 cats).

By "G-CSF" is meant one of the hematopoietic growth factors known as granulocyte colony stimulating factors. The biological activities of G-CSFs include:

10 stimulating the differentiation of a small number of progenitor "stem cells" into the variety of blood cell lines, stimulating the proliferation of those blood cell lines and stimulating the ultimate differentiation of mature blood cells from those lines. The preferred

15 G-CSF polypeptides for treating or preventing infections in canine or feline animals are human and canine, and may be naturally-derived or the product of genetically engineered host cells containing a DNA sequence encoding G-CSF.

20 The DNA encoding the G-CSF gene is a genomic DNA sequence, a cDNA sequence or a manufactured (or synthetic) DNA sequence which is expressed in a prokaryotic or eukaryotic host cell as a polypeptide having part or all of the primary structural

25 conformation and the hematopoietic biological properties of naturally-occurring G-CSF. A biologically functional plasmid or viral DNA vector containing a DNA sequence encoding G-CSF may be used to transform or transfect a prokaryotic or eukaryotic host cell to produce cell

30 lines expressing the G-CSF polypeptide, glycosylated or unglycosylated.

The various forms of G-CSF, including their preparation and purification, useful in a method for treating or preventing infections in canine or feline

35 animals commonly owned are described in detail in U.S. Patent 4,810,643, which is hereby incorporated by

- 9 -

reference. U.S. 4,810,643 describes and claims novel gene segments, biologically functional recombinant plasmids and viral DNA vectors and prokaryotic and eukaryotic host cells, which contain a G-CSF gene or a  
5 genetically engineered variant of a G-CSF gene. The host cells express biologically active G-CSF or a genetically engineered variant of G-CSF.

This application describes the isolation and characterization of a canine G-CSF gene and in  
10 particular describes and claims novel gene segments, biologically functional recombinant plasmids and viral DNA vectors, and prokaryotic and eukaryotic host cells, which contain a canine G-CSF gene or a genetically engineered variant of a canine G-CSF gene. The host  
15 cells transformed or transfected with the recombinant plasmids or viral DNA vectors express biologically active G-CSF or a genetically engineered variant of G-CSF. The protein expressed is purified using methods known to those skilled in the art.

20 DNA sequences coding for all or a part of G-CSF having the canine amino acid sequence are provided. Such DNA sequences include: the incorporation of codons "preferred" for expression by selected non-mammalian hosts (e.g. E. coli preferred  
25 codons, see Nucleic Acids Res. 1986 vol.14 (13) pp 5125-5143); the provision of sites for cleavage by restriction endonuclease enzymes; the provision of DNA sequences which reduce or eliminate secondary structure interactions which inhibit transcription and/or  
30 translation; and the provision of additional initial, terminal or intermediate DNA sequences which facilitate incorporation into expression vectors. The DNA sequences of the invention also include sequences having an optimized ribosome binding site, and sequences which  
35 enhance transcription, translation, and/or secretion of the protein product.

- 10 -

The present invention also provides DNA sequences coding for expression of polypeptide analogs or derivatives of canine G-CSF which differ from naturally-occurring forms in terms of the identity or location of one or more amino acid residues (i.e., deletion analogs containing less than all of the residues specified for canine G-CSF; substitution analogs, wherein one or more residues specified are replaced by other residues; and in addition, analogs wherein one or more amino acid residues are added to a terminal or medial portion of the polypeptide) and which share the properties of naturally-occurring forms.

Also comprehended by the present invention is that class of polypeptide coded for by portions of the DNA complement to the top strand canine cDNA of Figure 2, i.e., "complementary inverted proteins" as described by Tramontano, et al., Nucleic Acids Res., 12, 5049-5059 (1984).

The present invention relates to purified and isolated polypeptide products having part or all of the primary structural conformation (i.e., continuous sequence of amino acid residues) and the biological properties (e.g., immunological properties and in vitro biological activity) of naturally-occurring canine G-CSF including allelic variants thereof. These polypeptides are also characterized by being the product of chemical synthetic procedures or of procaryotic or eukaryotic host expression (e.g., by bacterial, yeast, higher plant, insect and mammalian cells (e.g. CHO or COS) in culture) of exogenous DNA sequences obtained by genomic or cDNA cloning or by gene synthesis. The products of typical yeast (e.g., Saccaromyces cerevisiae) or prokaryote [e.g., [Escherichia coli (E. coli)] host cells are free of association with any mammalian proteins. Depending upon the host employed, polypeptide of the invention is glycosylated with mammalian or other

- 11 -

eukaryotic carbohydrates or is non-glycosylated. Polypeptides of the invention may also include an initial methionine amino acid residue (at position -1).

In addition to the recombinant versions of

5 naturally-occurring allelic forms of canine G-CSF, the present invention also embraces other G-CSF products such as polypeptide analogs of canine G-CSF and fragments of canine G-CSF. All such forms of canine G-CSF may be useful in the method for treating or

10 preventing infections in canine or feline animals. Following the procedures of the published application by Alton, et al. (WO/83/04053), hereby incorporated by reference, one can readily design and manufacture genes coding for microbial expression of polypeptides having

15 primary conformations which differ from that herein specified for, in terms of the identity or location of one or more residues (e.g., substitutions, terminal and intermediate additions and deletions). Alternately, modifications of genomic and cDNA genes are readily

20 accomplished by well-known site-directed mutagenesis techniques which generate analogs and derivatives of canine G-CSF. Such products share the hematopoietic biological properties of canine G-CSF. As examples, products of the invention include those which are

25 foreshortened (e.g., by deletions); or those which are more stable to hydrolysis (and, therefore, have more pronounced or longer lasting effects than naturally-occurring); or which have been altered to delete one (or more) potential site(s) for n-linked or o-linked

30 glycosylation (which result in higher activities for yeast-produced products); or which have one or more cysteine residues deleted or replaced (for example, by alanine or serine residues) and are more easily isolated in active form from microbial systems; or which have one

35 or more tyrosine residues replaced by phenylalanine and bind more or less readily to G-CSF receptors on target

- 12 -

cells. Also comprehended are polypeptide fragments duplicating only part of the continuous amino acid sequence or secondary conformations of canine G-CSF.

According to another aspect of the present invention, the DNA sequence described herein which encodes G-CSF polypeptides is valuable for the information which it provides concerning the amino acid sequence of this canine protein (and similar mammalian proteins) which has heretofore been unavailable. The DNA sequences are also valuable as products useful in effecting the large scale microbial synthesis of G-CSF having the same amino acid sequence as canine G-CSF, by a variety of recombinant techniques. Put another way, DNA sequences provided by the invention are useful in generating new, and useful viral and plasmid DNA vectors, new and useful transformed and transfected prokaryotic and eukaryotic host cells (including bacterial, yeast, and mammalian cells grown in culture), and new and useful methods for cultured growth of such microbial host cells capable of expression of G-CSF having the canine amino acid sequence, variants or analogs. DNA sequences of the invention are also suitable materials for use as labelled probes in isolating canine G-CSF and related protein encoding cDNA and genomic DNA sequences of other mammalian species. DNA sequences are also useful in various alternative methods of protein synthesis (e.g., in insect cells) or in genetic therapy in mammals. DNA sequences of the invention are useful in developing transgenic mammalian species which may serve as eukaryotic "hosts" for production of G-CSF and G-CSF products in quantity. (See generally Palmiter, et al., Science, 22(4625), 809-814 (1983)).

Of applicability to canine G-CSF fragments and polypeptide analogs of the invention are reports of the immunological activity of synthetic peptides which substantially duplicate the amino acid sequence extant

- 13 -

in naturally-occurring proteins, glycoproteins and nucleoproteins. More specifically, relatively low molecular weight polypeptides have been shown to participate in immune reactions which are similar in duration and extent to the immune reactions of physiologically significant proteins such as viral antigens, polypeptide hormones, and the like. Included among the immune reactions of such polypeptides is the provocation of the formation of specific antibodies in immunologically active animals. (See, e.g., Lerner, et al., Cell, 23: 309-310 (1981); Ross, et al., Nature, 294: 654-656 (1981); Walter, et al., Proc. Natl. Acad. Sci. (USA), 77: 5197-5200 (1980); Lerner, et al., Proc. Natl. Acad. Sci. (USA), 78: 4882-4886 (1981); Wong, et al., Proc. Natl. Acad. Sci. (USA), 78: 7412-7416 (1981); Green, et al., Cell, 28: 477-587 (1982); Nigg, et al., Proc. Natl. Acad. Sci. (USA), 79: 5322-5326 (1982); Baron, et al., Cell, 28: 395-404 (1982); Dreesman, et al., Nature, 295: 183-190 (1982); and Lerner, Scientific American, 248 (2): 66-74 (1983)). See, also, Kaiser, et al. Science, 223: 249-255 (1984) relating to biological and immunological activities of synthetic peptides which approximately share secondary structures of peptide hormones but may not share their primary structural conformation.

All of the above mentioned forms, fragments, variants and analogs of canine G-CSF may be useful in the method of treating or preventing infections in canine or feline animals as described herein.

In another embodiment of the invention, one or more additional colony stimulating factors are administered to the infected animal with G-CSF, egs. GM-CSF, M-CSF, multi-CSF (IL-3). The CSFs are administered together or separately. In a further embodiment, animal infections are treated by administering G-CSF with one or more of: the interferons

- 14 -

(advantageously  $\alpha$ -interferon), IL-2, IL-6 and TNF or with a traditional antibiotic.

This application also describes pharmaceutical compositions of G-CSF having the canine amino acid sequence in a pharmaceutically acceptable carrier. These compositions may be administered intravascularly, intraperitoneally, subcutaneously, intramuscularly, or by infusion using forms known to the pharmaceutical art. For intravascular, intraperitoneal, subcutaneous, or intramuscular administration, active drug components may be combined with a suitable carrier such as water, saline, aqueous dextrose, and the like. Regardless of the route of administration selected, the compositions of the present invention are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those skilled in the art. An advantageous formulation is disclosed in commonly owned Ser. No. 285,159, hereby incorporated by reference. In one embodiment, sustained release formulations are used.

In one embodiment of the invention, G-CSF treatment is used in a prophylactic manner. For example, dogs or cats are treated prior to occurrences which may debilitate them, in order to boost and prime their capacity to fight off infections. Administration of the G-CSF can be made at the time the dogs or cats undergo surgery or radiation, etc.

Several variables will be taken into account by the ordinary artisan in determining the concentration of G-CSF in the therapeutic formulations and dosages to be administered. Variables include administration route and condition of the animal.

The following examples are presented by way of illustration of the invention and are specifically directed to procedures carried out prior to identification of canine G-CSF genomic and cDNA clones, to procedures resulting in such identification, and to



- 15 -

the sequencing, development of expression systems based on genomic, cDNA and manufactured (or synthetic) genes and verification of expression of G-CSF having the canine amino acid sequence, and analog products in such systems. The method of isolating the canine G-CSF gene described below can also be used to isolate other animal G-CSF genes, which in turn can be used in producing other animal G-CSFs. In addition, the examples illustrate methods for treating or preventing infections in canine animals, comprising administering an effective amount of G-CSF.

#### EXAMPLE 1

##### 15     Screening a Genomic Library for the canine G-CSF Gene

In this example, oligonucleotide probes were used to screen for a genomic clone containing a canine G-CSF gene. A phage (EMBL-3) canine genomic library was obtained from Clontech, plated out on E. coli strain NM538, and screened using <sup>32</sup>P phosphorylated oligonucleotide probes of the following sequences:

1. TCC CTG CCC CAG AGC TTC CTG CTC AAG TGC TTA GAG CAA GTG AGG AAG  
25   ATC CAG, and

2. GCC ATG CCG GCC TTC ACT TCT GCC TTC CAG CGC CGG GCA GGA GGG GTC  
CTG

30   A total of approximately  $1.0 \times 10^6$  phage were plated on 8 22 cm square petri dishes and plaque lifted in duplicate onto Gene Screen Plus transfer hybridization membranes. One set of membranes was hybridized to probe 1 and the other set was hybridized to probe 2  
35   using the procedures described in Maniatis et al., Molecular Cloning, A Laboratory Manual (Cold Spring

- 16 -

Harbor Laboratory, New York, 1982). Hybridizations were done at 55°C overnight in 6XSSC, 5X Denhardt's, 50 µg/ml sheared herring sperm DNA. A total of 1 positive clone was observed which hybridized to both probes. This clone was rescreened until an isolated plaque was obtained and was grown in a 3 liter culture and phage DNA was prepared as described in Maniatus, supra. This DNA was mapped by restriction enzyme digestion and Southern blotting using the radiolabeled probes. The mapping results showed that a Asp718 fragment of about 3700 bases contained the entire G-CSF region. DNA was digested with Asp718 to release an approximately 3700 bp canine G-CSF containing fragment which was subsequently subcloned into pUC19 at the Asp 718 site and further mapped by restriction endonuclease digests and Southern blotting.

A restriction endonuclease map (approximately 3.7 kb) of genomic DNA containing the canine G-CSF gene is shown in Figure 1. The sequence for the entire coding region of the mature canine G-CSF was determined by subcloning fragments into M13 and sequencing them by the dideoxy method described in Sanger et al., Proc. Natl. Acad. Sci. U.S.A. 74, 5463-5467 (1977). Sequences were confirmed or extended by utilizing internal primers off of the same clones. The sequence for the coding region was deduced by direct comparison with the human genomic G-CSF sequence (Figure 3) and is shown in Figure 2. Splice juncture sites and amino terminal processing of the protein were assumed to occur at the same places as the human G-CSF. The DNA sequence codes for a mature protein of the same length as the human G-CSF (174 amino acids) and the proteins are 81% homologous (see Figure 7).

- 17 -

EXAMPLE 2**Construction of Synthetic Canine G-CSF Genes  
and Expression of Such G-CSF Genes**

5

This example relates to preparation of manufactured genes encoding canine G-CSF and including E. coli preference codons, and to expression of such G-CSF.

10

Synthetic genes were designed to allow for the expression of canine granulocyte colony stimulating factor in E. coli [cG-CSF dna3 (Figures 4-6)]. Canine G-CSF is 174 amino acids in length and is 81% homologous to the human form of G-CSF (174 a.a.).

15

The gene cG-CSF dna3 (Figures 4-6) was designed with maximum bias for E. coli codon preference. For gene cG-CSF dna3, in addition to the coding sequence, an initiation ATG, leader and terminator sequences and 5' XbaI and 3' BamHI restriction sites were entered. The gene, cG-CSF dna3, was also designed to have minimum secondary interactions and sufficient unique restriction sites for subunit assembly and gene manipulation. BamHI and PstI sites were incorporated at positions identical to those found in the human G-CSF gene noted in commonly owned U.S. Patent 4,810,643. This allows for generation of unique human/canine hybrid genes and their protein products.

20

25

The gene was designed as two subunits (Subunit I (XbaI-HindIII), and Subunit II (HindIII-BamHI) for cloning into sequencing/expression vectors (Figure 6). Subunit I contains a short leader sequence with an XbaI cloning end and the ribosome binding site (RBS). Subunit II contains a pair of redundant stop codons and the BamHI cloning end.

30

35

Briefly stated, the protocol employed was generally as set out in the disclosure of co-owned

- 18 -

Alton, et al., PCT Publication No. WO83/04053, which is incorporated by reference herein. The gene was designed for initial assembly of component oligonucleotides into multiple duplexes which, in turn, were assembled into two discrete sections (Figure 6). These sections were designed for ready amplification and, upon removal from the amplification system, could be assembled sequentially or through a multiple fragment ligation into a suitable expression vector.

10 The construction of Sections I and II is illustrated in Figures 5 and 6. In the construction of Section I, as shown in Figures 5 and 6, 16 oligonucleotides were assembled into 8 duplexes. The 8 duplexes were then ligated to form Section I. It may also be noted in Figure 6 that Section I includes an upstream XbaI sticky end and a downstream HindIII sticky end useful for ligation to amplification and expression vectors and for ligation to Section II.

Section II was constructed as shown in Figures 5 and 6. For this construction, 16 oligonucleotides were assembled into 8 duplexes. The 8 duplexes were then ligated to form Section II as depicted in Figure 6. As also shown in Figure 6, Section II includes an upstream HindIII sticky end and a downstream BamHI sticky end useful for ligating into amplification and expression vectors, and to Section I.

Although any suitable vector may be employed to express this DNA, the expression plasmid pCFM536 may readily be used. This plasmid is described in U.S. Patent No. 4,710,473 hereby incorporated by reference. Control of expression in the pCFM536 plasmid is by means of a lambda pL promoter, which itself may be under the control of a CI857 repressor gene (such as is provided in E. coli strain FM5 (ATCC deposit 53911)).

35 Section I was initially cloned into M13 from XbaI to HindIII and sequenced by the dideoxy method

- 19 -

(Sanger supra). Section II was cloned into M13 from HindIII to EcoRI and was also sequenced by the dideoxy method. Section I was cut out of M13 from XbaI to HindIII and Section II was cut out of M13 from HindIII to EcoRI. These two fragments were then ligated with pCFM536 cut from XbaI to BamHI and transformed into E. coli strain FM5 to generate pCFM536cG-CSF.

This plasmid contains the  $\lambda$ pL promoter/operator region and has a temperature sensitive replicon. When E. coli strain FM5 harboring pCFM536cG-CSF is cultured at 28°C, the plasmid copy number is maintained at 10-20 copies/cell, and transcription from the  $\lambda$ pL promoter is regulated by a temperature sensitive repressor. Growth at 42°C results in an increased copy number and release of repression at the  $\lambda$ pL promoter. Recombinant G-CSF having the canine sequence begins to accumulate at elevated temperatures as the result of promoter activation and plasmid amplification. The  $\lambda$ pL promoter lies just upstream from the ribosome binding site and the methionine initiation codon of canine G-CSF. The transcription terminator, t-oop, lies just downstream from the two translational stop codons near the 3' end of the gene. Strain FM5 harboring the plasmid, pCFM536cG-CSF, expresses recombinant G-CSF having the canine sequence at up to 30% of the total cellular protein.

### EXAMPLE 3

#### 30 Construction of Canine G-CSF Analogs

This example relates to the use of recombinant methods to generate an analog of canine G-CSF wherein the cysteine at position 17 was individually replaced by a serine.

- 20 -

Site directed mutagenesis procedures according to Souza, et al., published PCT Application No. WO85/00817, published February 28, 1985, hereby incorporated by reference, were carried out using the  
5 oligonucleotide CTG CTG AAA TCC CTC GAG CAG.

#### EXAMPLE 4

##### E. coli Canine G-CSF Purification

10

The general purification method is disclosed in commonly owned Ser. No. 348,011 hereby incorporated by reference.

##### 15 Cell Breakage and Sarkosyl Solubilization and Oxidation

About 200 grams of cell paste were weighed out in 1.5 liters of cold water. The cell paste was dispersed with a homogenizer until completely  
20 dispersed. The homogenate was then passed through a Gaulin Homogenizer four times at 8000 psig. The material was then centrifuged in the Beckman J2 21 centrifuge using the JA 10 rotor at 9000 rpm for 30 minutes at 4°C. The supernatant was decanted and  
25 discarded. The pellet was resuspended in 1.5 liters of cold water and again centrifuged in the Beckman J2 21 centrifuge using the JA 10 rotor at 9000 rpm for 30 minutes at 4°C. The supernatant was decanted and discarded. The pellet was resuspended in 760 mL water  
30 and 40 mL 1M Tris, pH 8.0 was added followed by 200 mL 10% Sarkosyl. After this material stirred at room temperature for about ten minutes, 1 mL 1% copper sulfate pentahydrate was added. This material was stirred at room temperature overnight (approximately  
35 16 hours). The material was then centrifuged in the Beckman J2 21 centrifuge using the JA 10 rotor at 9000

- 21 -

rpm for 30 minutes at 4°C. The supernatant was decanted and saved. The pellets were discarded.

#### Dowex Removal of Sarkosyl

5

To the supernatant was added 1 liter of cold water and then 2 liters cold 20 mM Tris, pH 8.0 and then 800 grams prepared Dowex (see Ser. No. 348,011 hereby incorporated by reference) was added. This slurry was  
10 stirred at 4°C for 90 minutes. The slurry was poured through a column and the flow through collected. The resin was washed with 800 mL cold 20 mM Tris, pH 8.0 which was added to the flow through giving 4800 mL.

#### 15 DE52 Cellulose Ion Exchange Chromatography

About 4800 mL of material was loaded directly onto a 200 mL DE52 cellulose ion exchange column equilibrated in 20 mM Tris, pH 8.0. The product was  
20 eluted off of the column using 100 mM NaCl in 20 mM Tris, pH 9.0. About 1270 mL was collected at approximately 0.8 mg/mL, giving approximately 1 gram.

#### CM-Sepharose Fast Flow Chromatography

25

The DE52 100 mM NaCl material was concentrated using a Pellicon system (with a 10,000 MW membrane) to approximately 200 mL. The material was adjusted to pH 5.4 using 50% acetic acid. Six volumes of cold water  
30 were added and the material was then loaded directly onto a 50 mL CM-Sepharose Fast Flow ion exchange column equilibrated in 20 mM sodium acetate, pH 5.4. The product was eluted off of the column using a 1 liter gradient from 0-0.2 M NaCl in 20 mM sodium acetate,  
35 pH 5.4. About 100 10 mL fractions were collected. Based on the chromatogram results the fractions of

- 22 -

interest were analyzed on a 15% SDS gel. Based on the gel results, fractions 30-51 were pooled giving 258 mL at approximately 2.6 mg/mL, or 685 mgs.

## 5 Diafiltration

The CM pool was adjusted to pH 3.5 using 0.1 N HCL and then diafiltered using a Pellicon with a 10,000 MW membrane vs. 0.35 mM HCL-Water. The final volume was  
10 adjusted to 685 mL to give material at a final concentration of 1 mg/mL.

## EXAMPLE 5

15

### In/vivo Activity of Canine G-CSF

Two young adult, healthy mixed breed dogs (one 25 kg male, one 28.6 kg female) were used for this study. The dogs were acclimated to the hospital  
20 environment for one week prior to the onset of the study. Complete blood and platelet counts were done three days prior and then immediately prior to the first injection of recombinant cG-CSF. Recombinant E. coli G-CSF having the amino acid sequence of canine G-CSF was  
25 diluted in sterile water to 100ug/ml and placed in multiple dose vials. The G-CSF was maintained at 4°C.

A dosage of 5 ug/kg/day was administered subcutaneously to each dog for 4 weeks at the same time each day. Blood for a CBC and platelet count was drawn  
30 immediately prior to each G-CSF injection and submitted to the clinical pathology laboratory for evaluation. Daily blood counts were performed until three consecutive daily counts remained stable. Blood was then drawn every other day for two weeks, then ev ry  
35 third day th final week.



- 23 -

After 28 days, G-CSF administration was discontinued. Blood counts were followed every other day to determine how rapidly they returned to normal. Once within normal range, G-CSF was started again at the same dosage and administered for another five days to determine the leukocyte response.

Physical examinations were performed on a daily basis. Karnofsky's performance scores were assigned daily to both animals. Body weights and body temperatures were recorded daily. In addition, toxicity evaluation was performed daily. The mean white blood cell count prior to administration of G-CSF was 8,650/ul (neutrophils: 4,880/ul; lymphocytes: 2,398/ul; monocytes: 667/ul; eosinophils: 704/ul; and platelets: 297,000/ul). Twenty-four hours following the first injection of G-CSF, the mean white blood cell count was 39,150/ul (neutrophils: 31,257/ul; neutrophilic bands: 391/ul; lymphocytes: 2,803/ul; monocytes: 2,951/ul; eosinophils: 1,747/ul; platelets: 322,500/ul). This represents a 4.5 fold increase in total white blood cell count within 24 hours. Neutrophils increased by a factor of 6.4 (see Figure 8). Monocytes rose by a factor of 4.4 (see Figure 9). Although the dosage of G-CSF remained at 5 ug/kg/day, an additional increase in blood counts was noted on day eleven. Mean white blood cell count on day nine was 32,550/ul (mean neutrophil count: 26,682/ul). On day eleven, the mean white blood cell count was 69,200/ul (mean neutrophil count: 58,764/ul) representing an additional two-fold increase from day nine to day eleven and an eight-fold increase from day one (prior to G-CSF administration). Blood counts remained elevated throughout the 28 day period of administration of G-CSF in one dog. In the second dog there were 3 days on which decreases in the leukocyte counts were evident 24 hours after administration of a reduced dosage. Counts returned to their pretreatment

- 24 -

levels by the fifth day after G-CSF was stopped. Upon resumption of G-CSF administration, the mean white blood cell count increased by a factor of 6.3 (from mean of 9,450/ul to mean of 59,500/ul). These elevated counts  
5 persisted until G-CSF administration was discontinued five days later (See Figures 8 and 9).

Recombinant G-CSF having the amino acid sequence of canine G-CSF increased leukocyte counts (primarily neutrophils) and leukocyte counts were  
10 maintained at elevated levels as long as administration of the G-CSF was continued. Initial increases in leukocyte counts were most likely due to demargination of blood cells. The decrease in leukocyte counts observed following a reduced G-CSF dosage followed by a  
15 rapid return to elevated leukocyte levels with a full dosage demonstrate a rapid, dose-dependent response. There was no development of neutralizing antibodies to the G-CSF.

20

\* \* \*

While the present invention has been described  
25 in terms of preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations which come within the scope of the invention as claimed.

30

35

- 25 -

## WHAT IS CLAIMED IS:

1. A purified and isolated polypeptide having part or all of the primary structural conformation and  
5 the biological properties of naturally-occurring canine granulocyte colony stimulating factor and characterized by being the product of prokaryotic or eukaryotic expression of an exogenous DNA sequence.
- 10 2. A polypeptide according to claim 1 wherein the exogenous DNA sequence is a genomic DNA sequence.
3. A polypeptide according to claim 1 wherein the exogenous DNA sequence is a cDNA sequence.
- 15 4. A polypeptide according to claim 1 wherein the exogenous DNA sequence is a manufactured DNA sequence.
- 20 5. A polypeptide according to claim 1 wherein the exogenous DNA sequence is carried on an autonomously replicating DNA plasmid or viral vector.
6. A polypeptide according to claim 1  
25 possessing part or all of the primary structural conformation of canine granulocyte colony stimulating factor as set forth in Figure 2 or any naturally occurring allelic variant thereof.
- 30 7. A polypeptide according to claim 1 which has the immunological properties of naturally-occurring canine granulocyte colony stimulating factor.
8. A polypeptide according to claim 1 which  
35 has the in vitro biological activity of naturally-occurring canine granulocyte colony stimulating factor.

- 26 -

9. A DNA sequence for use in securing expression in a prokaryotic or eukaryotic host cell of a polypeptide product having at least a part of the primary structural conformation and the biological properties of naturally-occurring canine granulocyte colony stimulating factor, said DNA sequence set out in Figure 2 or its complimentary strand.

10. A prokaryotic or eukaryotic host cell transformed or transfected with a DNA sequence according to claim 9 in a manner allowing the host cell to express the polypeptide product.

11. A polypeptide product of the expression of a DNA sequence according to claim 9 in a prokaryotic or eukaryotic host cell.

12. A polypeptide product according to claim 11 wherein the polypeptide product is glycosylated or unglycosylated.

13. A purified and isolated DNA sequence coding for prokaryotic or eukaryotic host cell expression of a polypeptide having part or all of the primary structural conformation and the biological properties of canine granulocyte colony stimulating factor.

14. A genomic DNA sequence according to claim 13.

15. A cDNA sequence according to claim 13.

16. A DNA sequence according to claim 13 and including one or more codons preferred for expression in E. coli cells.

- 27 -

17. A DNA sequence according to claim 13 and including one or more codons preferred for expression in yeast cells.

5           18. A DNA sequence coding for a polypeptide fragment or polypeptide analog of naturally-occurring canine granulocyte colony stimulating factor.

10           19. A biologically functional plasmid or viral DNA vector containing a DNA sequence according to claim 9.

15           20. A prokaryotic or eukaryotic host cell stably transformed or transfected with the biologically functional plasmid or viral DNA vector according to claim 19.

20           21. A biologically functional plasmid or viral DNA vector containing a DNA sequence according to claim 13.

25           22. A prokaryotic or eukaryotic host cell stably transformed or transfected with the biologically functional plasmid or viral DNA vector according to claim 21.

30           23. A biologically functional plasmid or viral DNA vector containing a DNA sequence according to claim 18.

35           24. A prokaryotic or eukaryotic host cell stably transformed or transfected with the biologically functional plasmid or viral DNA vector according to claim 23.

- 28 -

25. A polypeptide product of the expression in a prokaryotic or eukaryotic host cell of a DNA sequence according to claims 13 or 18.

5           26. A synthetic polypeptide having part or all of the amino acid sequence as set forth in Figure 2 and having the in vitro biological activities of naturally-occurring canine granulocyte colony stimulating factor.

10

27. A synthetic polypeptide having part or all of the secondary conformation and part or all of the amino acid sequence set forth in Figure 2 and having the biological properties of naturally-occurring canine  
15 granulocyte colony stimulating factor.

28. A process for the production of a polypeptide having part or all of the primary structural conformation and the biological properties of naturally-  
20 occurring canine granulocyte colony stimulating factor, the process comprising: growing, under suitable nutrient conditions, prokaryotic or eukaryotic host cells transformed or transfected with a biologically functional plasmid or viral DNA vector according to  
25 claims 19, 21, or 23, and isolating desired polypeptide products of the expression of DNA sequences in the biologically functional plasmid or viral DNA vector.

29. A method for treating or preventing  
30 infections in a canine or feline animal comprising administering a composition which comprises a therapeutically effective treating amount or preventive amount of granulocyte colony stimulating factor.

35

- 29 -

30. A method according to claim 29 wherein said administering step comprises administering said granulocyte colony stimulating factor and one or more compounds selected from the group consisting of:

5 GM-CSF, M-CSF, IL-3, interferon, IL-2, IL-6, TNF and a traditional antibiotic.

31. A method according to claim 29 wherein the composition is administered by the parenteral route.

10

32. A method according to claim 29 wherein the animal is a dog or cat.

33. A method according to claim 29 wherein  
15 the composition is a granulocyte colony stimulating factor having the human amino acid sequence.

34. A method according to claim 33 wherein the granulocyte colony stimulating factor is naturally-  
20 derived or is derived from genetically engineered host cells containing a genomic DNA sequence, a cDNA sequence or a manufactured DNA sequence encoding human granulocyte colony stimulating factor.

25 35. A method according to claim 29 wherein the composition is a granulocyte colony stimulating factor having the canine amino acid sequence.

36. A method according to claim 35 wherein  
30 the granulocyte colony stimulating factor is naturally-derived or is derived from genetically engineered host cells containing a genomic DNA sequence, a cDNA sequence or a manufactured DNA sequence encoding canine granulocyte colony stimulating factor.

35

- 30 -

37. A method according to claim 35 wherein the granulocyte colony stimulating factor is the polypeptide product of the expression in a prokaryotic or eukaryotic host cell of DNA sequence according to  
5 claims 9, 13, or 18.

38. A method for treating cancer in canine or feline animals comprising administering a composition which comprises a therapeutically effective treating or  
10 preventing amount of granulocyte colony stimulating factor in conjunction with chemotherapy.

39. A pharmaceutical composition for treating infections in canine or feline animals comprising a  
15 therapeutically effective amount of a granulocyte colony stimulating factor, in a pharmaceutically acceptable carrier.

20

25


30

35



1 / 16

**Restriction Map of  
Canine G-CSF**



**FIG. 1**

2 / 16

Ala Pro Leu Gly Pro Thr Gly Pro Leu Pro Gln Ser Phe Leu Leu  
gcc ccc ctg ggc cct acc ggc ccc ctg ccc cag agc ttc ctg ctg

Lys Cys Leu Glu Gln Met Arg Lys Val Gln Ala Asp Gly Thr Ala  
aag tgc cta gag caa atg agg aag gtc cag gct gat ggc acg gcg

Leu Gln Glu Thr Leu Cys Ala Thr His Gln Leu Cys His Pro Glu  
ctg cag gag acg ctg tgt gcc acc cac cag ctg tgc cat cct gag

Glu Leu Val Leu Leu Gly His Ala Leu Gly Ile Pro Gln Pro Pro  
gag ttg gtg ctg ctc ggc cac gct ctg ggc atc ccc cag cct ccc

Leu Ser Ser Cys Ser Ser Gln Ala Leu Gln Leu Met Gly Cys Leu  
ctg agc agc tgc tcc agc agc ggc ctg cag ctg atg ggc tgc ctg

Arg Gln Leu His Ser Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln  
cgt caa ctc cac agc ggc ctc ttc ctc tac cag ggc ctc ctg cag

FIG. 2-A

SUBSTITUTE SHEET

3 / 16

Ala Leu Ala Gly Ile Ser Pro Glu Leu Ala Pro Thr Leu Asp Thr  
gcc ctg gca ggg ata tcc ccc gag tta gcg ccc acc ttg gac aca

Leu Gln Leu Asp Thr Thr Asp Phe Ala Ile Asn Ile Trp Gln Gln  
ctg cag ctg gac acc acc gac ttt gcc atc aac atc tgg cag cag

Met Glu Asp Leu Gly Met Ala Pro Ala Val Pro Thr Gln Gly  
atg gaa gat cta gga atg gcc ccc gcc gtg cca cct acc cag ggc

Thr Met Pro Ala Phe Thr Ser Ala Phe Gln Arg Arg Ala Gly Gly  
acc atg cca gcc ttc acc tcc gcc ttc cag cgc cgc gga ggt

Val Leu Val Ala Ser Asn Leu Gln Ser Phe Leu Glu Leu Ala Tyr  
gtc ctg gtg gcc tcc aac ctg cag agc ttc ctg gag ctg gca tat

Arg Ala Leu Arg His Phe Ala Lys Pro  
cgc gct ctg cgc cac ttt gcc aaa ccc

FIG. 2-B

SUBSTITUTE SHEET

4 / 16

GGGGACAGGCTTGAGAATCCCAAAGGAGAGGGGCAAGGACACTGCCCCCGCAAGTCTGCCAGAGCAGAGAGGGAGACCCCGACTCAGCTGCCACTTCCC 100  
 CACAGGCTCGTGCCGCTCCAGGGCTCTATCAGCGGCTCAGCCTTTGTTTCAGCTGTTCTGTTCAAACACTCTGGGGCCATTTCAGGCCCTGGGTGGGGCAGC 200  
 GGGAGGAAGGAGTTTGAGGGGGGCAAGGCGACGTCAAAGGAGGATCAGAGATTCACAACTTTCACAAACTTTCGCAAAACAGCTTTTGTTCCAACCCC 300  
 CCTGCATTGTCTTGGACACCAAAATTTCATAAATCTCTGGGAAGTTATTACTAAGCCTTAGTCGTGGCCCCCAGGTAATTTCTCTCCAGGCCCTCCATGGGGT 400  
 TATGTATAAAGGGCCCCCTAGAGCTGGGCCCCCAAAACAGCCCGGAGCCTGCAGCCCCAGCCCCACCCAGACCCCATGGCTGGACCTGCCACCCAGAGCCCCA 500  
 TGGTGAGTGTCTTG6CCCAGGATGGGAGAGCCGCCCTGCCCTGGCATGGGAGGGGCTGGTGTGACAGAGGGGCTGGGGATCCCCGTTCCT 600  
 GGGAAATGGGGATTAAAGGCACCCAGTGTCCTCCGAGAGGGCCTCAGGTGGTAGGGAACAGCATGTCCTCTGAGCCCCGCTCTGTCTCCCCAG 700  
 euleuTrpHisSerAlaLeuTrpThrValGlnGluAlaThrProLeuGlyProAlaSerSerLeuProGlnSerPheLeuLeuLysCysLeuGluGlnVa 800  
 TGCTGTGGCACAGTGCACCTCTGGACAGTGCAGGAAGCCACCCCTGGGCCCTGCCAGCTCCCTGCCCCAGAGCTTCCTGCTCAAGTGTAGAGCAAGT 900  
 TArgLysIleGlnGlyAspGlyAlaAlaLeuGlnGluLysLeu 35  
 GAGGAAGATCCAGGGCGATGGCGACGCTCCAGGAGAAGCTGTGAGTGAGGTGGGTGAGAGGGCTGTGGAGGGGAAGCCCCGGTGGGGAGAGCTAAGGGG

-20 -18  
 etLysLeuMeta  
 TGAAGCTGATGG

-30  
 MetAlaGlyProAlaThrGlnSerProM

-16  
 laLeuGlnLeuL  
 CCCTGCAGCTGC

-10 -1 +1 10 20 30 35

FIG. 3-A

5 / 16

GATGGAAC TGCAGG6CCAACA TCC TCTGGAAGGGACATGGGAGAATAT TAGGAGCAGTGGAGCTG6GGAAGGCTGGGAAGGACT TGGGGAGGAGGACCT 1000  
 TGGTGGGACAGTGTCTCGGGAGGGCTGGCTGGGATGGGAGTGGAGGCATCACATTCAGGAGAAAGGGCAAGGGCCCTGTGTGAGATCAGAGAGTGGGGTG 1100  
 CAGGGCAGAGAGGAAC TGAACAGCC TGGCAGGACATGGAGGGAGGGGAAAGACCAGAGAGTGGGGAGGACCCGGGAAGGAGCGGCCGACCGGC 1200  
 GAGTCTCACTCAGCATCCTTCCATCCCCAGTGTGCCACCTACAAGCTGTGCCACCCCGAGGAGCTGGTGTCTCGGACACACTCTCTGGGCATCCCCCTGGG 1300  
 CysAlaThrTyrLysLeuCysHisProGluGluLeuValLeuLeuGlyHisSerLeuGlyIleProTrpA 50  
 36 40  
 60 70 71  
 laProLeuSerSerCysProSerGlnAlaLeuGlnLeu 72  
 CTCCCTGAGCAGCTGCCAGCCAGGCCCTGCAGCTGGTGTGAGTGTGAGGAAAGGATAAGGCTAATGAGGAGGGGGAAGGAGGAGGAAACACCCATGGG 1400  
 CTCCCCCATGTCTCCAGGTCCCAAGCTGGGGGCTGACGTATCTCAGGCAGCACCCCTAACTCTTCCGCTCTGTCTCACAGG 1500  
 AlaGlyCysLeuSerGln 72  
 80 90 100 110  
 LeuHisSerGlyLeuPheLeuTyrGlnGlyLeuLeuGlnAlaLeuGluGlyIleSerProGluLeuGlyProThrLeuAspThrLeuGlnLeuAspValA 1600  
 CTCCATAGCGGCTTTTCCCTCTACCCAGGGCTCTCTGCAGGCCCTGGAAAGGATCTCCCCCGAGTGGGTCCACCTTGGACACACTGCAGCTGGACGTCG 1700  
 120  
 laAspPheAlaThrThrIleTrpGlnGln  
 CCGACTTTGCCACCACCATCTGGCAGCAGGTGAGCCTTGTGGGCAGGGTGGCCCAAGGTGCTGTGGCATCTCTGGGCACCACAGCCGGGCTGTGTATGG

FIG. 3-B

6 / 16

121 MetGluG  
 1800  
 1900  
 2000  
 2100  
 2200  
 2300  
 2400  
 2500  
 2600  
 2700  
 2800  
 2900  
 3000  
 3070

GCCC.TGTCATGCTGTCAGCCCCCAGCATTTCCCTCATTTTGTAATAACGCCCACTCAGAGGGCCCAACCACTGATCACAGCTTTCCCCCACAGATGGAAG  
 130  
 140  
 150  
 160  
 170  
 174  
 177  
 180  
 183  
 186  
 189  
 192  
 195  
 198  
 201  
 204  
 207  
 210  
 213  
 216  
 219  
 222  
 225  
 228  
 231  
 234  
 237  
 240  
 243  
 246  
 249  
 252  
 255  
 258  
 261  
 264  
 267  
 270  
 273  
 276  
 279  
 282  
 285  
 288  
 291  
 294  
 297  
 300  
 303  
 306  
 309  
 312  
 315  
 318  
 321  
 324  
 327  
 330  
 333  
 336  
 339  
 342  
 345  
 348  
 351  
 354  
 357  
 360  
 363  
 366  
 369  
 372  
 375  
 378  
 381  
 384  
 387  
 390  
 393  
 396  
 399  
 402  
 405  
 408  
 411  
 414  
 417  
 420  
 423  
 426  
 429  
 432  
 435  
 438  
 441  
 444  
 447  
 450  
 453  
 456  
 459  
 462  
 465  
 468  
 471  
 474  
 477  
 480  
 483  
 486  
 489  
 492  
 495  
 498  
 501  
 504  
 507  
 510  
 513  
 516  
 519  
 522  
 525  
 528  
 531  
 534  
 537  
 540  
 543  
 546  
 549  
 552  
 555  
 558  
 561  
 564  
 567  
 570  
 573  
 576  
 579  
 582  
 585  
 588  
 591  
 594  
 597  
 600  
 603  
 606  
 609  
 612  
 615  
 618  
 621  
 624  
 627  
 630  
 633  
 636  
 639  
 642  
 645  
 648  
 651  
 654  
 657  
 660  
 663  
 666  
 669  
 672  
 675  
 678  
 681  
 684  
 687  
 690  
 693  
 696  
 699  
 702  
 705  
 708  
 711  
 714  
 717  
 720  
 723  
 726  
 729  
 732  
 735  
 738  
 741  
 744  
 747  
 750  
 753  
 756  
 759  
 762  
 765  
 768  
 771  
 774  
 777  
 780  
 783  
 786  
 789  
 792  
 795  
 798  
 801  
 804  
 807  
 810  
 813  
 816  
 819  
 822  
 825  
 828  
 831  
 834  
 837  
 840  
 843  
 846  
 849  
 852  
 855  
 858  
 861  
 864  
 867  
 870  
 873  
 876  
 879  
 882  
 885  
 888  
 891  
 894  
 897  
 900  
 903  
 906  
 909  
 912  
 915  
 918  
 921  
 924  
 927  
 930  
 933  
 936  
 939  
 942  
 945  
 948  
 951  
 954  
 957  
 960  
 963  
 966  
 969  
 972  
 975  
 978  
 981  
 984  
 987  
 990  
 993  
 996  
 999  
 1002  
 1005  
 1008  
 1011  
 1014  
 1017  
 1020  
 1023  
 1026  
 1029  
 1032  
 1035  
 1038  
 1041  
 1044  
 1047  
 1050  
 1053  
 1056  
 1059  
 1062  
 1065  
 1068  
 1071  
 1074  
 1077  
 1080  
 1083  
 1086  
 1089  
 1092  
 1095  
 1098  
 1101  
 1104  
 1107  
 1110  
 1113  
 1116  
 1119  
 1122  
 1125  
 1128  
 1131  
 1134  
 1137  
 1140  
 1143  
 1146  
 1149  
 1152  
 1155  
 1158  
 1161  
 1164  
 1167  
 1170  
 1173  
 1176  
 1179  
 1182  
 1185  
 1188  
 1191  
 1194  
 1197  
 1200  
 1203  
 1206  
 1209  
 1212  
 1215  
 1218  
 1221  
 1224  
 1227  
 1230  
 1233  
 1236  
 1239  
 1242  
 1245  
 1248  
 1251  
 1254  
 1257  
 1260  
 1263  
 1266  
 1269  
 1272  
 1275  
 1278  
 1281  
 1284  
 1287  
 1290  
 1293  
 1296  
 1299  
 1302  
 1305  
 1308  
 1311  
 1314  
 1317  
 1320  
 1323  
 1326  
 1329  
 1332  
 1335  
 1338  
 1341  
 1344  
 1347  
 1350  
 1353  
 1356  
 1359  
 1362  
 1365  
 1368  
 1371  
 1374  
 1377  
 1380  
 1383  
 1386  
 1389  
 1392  
 1395  
 1398  
 1401  
 1404  
 1407  
 1410  
 1413  
 1416  
 1419  
 1422  
 1425  
 1428  
 1431  
 1434  
 1437  
 1440  
 1443  
 1446  
 1449  
 1452  
 1455  
 1458  
 1461  
 1464  
 1467  
 1470  
 1473  
 1476  
 1479  
 1482  
 1485  
 1488  
 1491  
 1494  
 1497  
 1500  
 1503  
 1506  
 1509  
 1512  
 1515  
 1518  
 1521  
 1524  
 1527  
 1530  
 1533  
 1536  
 1539  
 1542  
 1545  
 1548  
 1551  
 1554  
 1557  
 1560  
 1563  
 1566  
 1569  
 1572  
 1575  
 1578  
 1581  
 1584  
 1587  
 1590  
 1593  
 1596  
 1599  
 1602  
 1605  
 1608  
 1611  
 1614  
 1617  
 1620  
 1623  
 1626  
 1629  
 1632  
 1635  
 1638  
 1641  
 1644  
 1647  
 1650  
 1653  
 1656  
 1659  
 1662  
 1665  
 1668  
 1671  
 1674  
 1677  
 1680  
 1683  
 1686  
 1689  
 1692  
 1695  
 1698  
 1701  
 1704  
 1707  
 1710  
 1713  
 1716  
 1719  
 1722  
 1725  
 1728  
 1731  
 1734  
 1737  
 1740  
 1743  
 1746  
 1749  
 1752  
 1755  
 1758  
 1761  
 1764  
 1767  
 1770  
 1773  
 1776  
 1779  
 1782  
 1785  
 1788  
 1791  
 1794  
 1797  
 1800  
 1803  
 1806  
 1809  
 1812  
 1815  
 1818  
 1821  
 1824  
 1827  
 1830  
 1833  
 1836  
 1839  
 1842  
 1845  
 1848  
 1851  
 1854  
 1857  
 1860  
 1863  
 1866  
 1869  
 1872  
 1875  
 1878  
 1881  
 1884  
 1887  
 1890  
 1893  
 1896  
 1899  
 1902  
 1905  
 1908  
 1911  
 1914  
 1917  
 1920  
 1923  
 1926  
 1929  
 1932  
 1935  
 1938  
 1941  
 1944  
 1947  
 1950  
 1953  
 1956  
 1959  
 1962  
 1965  
 1968  
 1971  
 1974  
 1977  
 1980  
 1983  
 1986  
 1989  
 1992  
 1995  
 1998  
 2001  
 2004  
 2007  
 2010  
 2013  
 2016  
 2019  
 2022  
 2025  
 2028  
 2031  
 2034  
 2037  
 2040  
 2043  
 2046  
 2049  
 2052  
 2055  
 2058  
 2061  
 2064  
 2067  
 2070  
 2073  
 2076  
 2079  
 2082  
 2085  
 2088  
 2091  
 2094  
 2097  
 2100  
 2103  
 2106  
 2109  
 2112  
 2115  
 2118  
 2121  
 2124  
 2127  
 2130  
 2133  
 2136  
 2139  
 2142  
 2145  
 2148  
 2151  
 2154  
 2157  
 2160  
 2163  
 2166  
 2169  
 2172  
 2175  
 2178  
 2181  
 2184  
 2187  
 2190  
 2193  
 2196  
 2199  
 2202  
 2205  
 2208  
 2211  
 2214  
 2217  
 2220  
 2223  
 2226  
 2229  
 2232  
 2235  
 2238  
 2241  
 2244  
 2247  
 2250  
 2253  
 2256  
 2259  
 2262  
 2265  
 2268  
 2271  
 2274  
 2277  
 2280  
 2283  
 2286  
 2289  
 2292  
 2295  
 2298  
 2301  
 2304  
 2307  
 2310  
 2313  
 2316  
 2319  
 2322  
 2325  
 2328  
 2331  
 2334  
 2337  
 2340  
 2343  
 2346  
 2349  
 2352  
 2355  
 2358  
 2361  
 2364  
 2367  
 2370  
 2373  
 2376  
 2379  
 2382  
 2385  
 2388  
 2391  
 2394  
 2397  
 2400  
 2403  
 2406  
 2409  
 2412  
 2415  
 2418  
 2421  
 2424  
 2427  
 2430  
 2433  
 2436  
 2439  
 2442  
 2445  
 2448  
 2451  
 2454  
 2457  
 2460  
 2463  
 2466  
 2469  
 2472  
 2475  
 2478  
 2481  
 2484  
 2487  
 2490  
 2493  
 2496  
 2499  
 2502  
 2505  
 2508  
 2511  
 2514  
 2517  
 2520  
 2523  
 2526  
 2529  
 2532  
 2535  
 2538  
 2541  
 2544  
 2547  
 2550  
 2553  
 2556  
 2559  
 2562  
 2565  
 2568  
 2571  
 2574  
 2577  
 2580  
 2583  
 2586  
 2589  
 2592  
 2595  
 2598  
 2601  
 2604  
 2607  
 2610  
 2613  
 2616  
 2619  
 2622  
 2625  
 2628  
 2631  
 2634  
 2637  
 2640  
 2643  
 2646  
 2649  
 2652  
 2655  
 2658  
 2661  
 2664  
 2667  
 2670  
 2673  
 2676  
 2679  
 2682  
 2685  
 2688  
 2691  
 2694  
 2697  
 2700  
 2703  
 2706  
 2709  
 2712  
 2715  
 2718  
 2721  
 2724  
 2727  
 2730  
 2733  
 2736  
 2739  
 2742  
 2745  
 2748  
 2751  
 2754  
 2757  
 2760  
 2763  
 2766  
 2769  
 2772  
 2775  
 2778  
 2781  
 2784  
 2787  
 2790  
 2793  
 2796  
 2799  
 2802  
 2805  
 2808  
 2811  
 2814  
 2817  
 2820  
 2823  
 2826  
 2829  
 2832  
 2835  
 2838  
 2841  
 2844  
 2847  
 2850  
 2853  
 2856  
 2859  
 2862  
 2865  
 2868  
 2871  
 2874  
 2877  
 2880  
 2883  
 2886  
 2889  
 2892  
 2895  
 2898  
 2901  
 2904  
 2907  
 2910  
 2913  
 2916  
 2919  
 2922  
 2925  
 2928  
 2931  
 2934  
 2937  
 2940  
 2943  
 2946  
 2949  
 2952  
 2955  
 2958  
 2961  
 2964  
 2967  
 2970  
 2973  
 2976  
 2979  
 2982  
 2985  
 2988  
 2991  
 2994  
 2997  
 3000  
 3003  
 3006  
 3009  
 3012  
 3015  
 3018  
 3021  
 3024  
 3027  
 3030  
 3033  
 3036  
 3039  
 3042  
 3045  
 3048  
 3051  
 3054  
 3057  
 3060  
 3063  
 3066  
 3069  
 3072  
 3075  
 3078  
 3081  
 3084  
 3087  
 3090  
 3093  
 3096  
 3099  
 3102  
 3105  
 3108  
 3111  
 3114  
 3117  
 3120  
 3123  
 3126  
 3129  
 3132  
 3135  
 3138  
 3141  
 3144  
 3147  
 3150  
 3153  
 3156  
 3159  
 3162  
 3165  
 3168  
 3171  
 3174  
 3177  
 3180  
 3183  
 3186  
 3189  
 3192  
 3195  
 3198  
 3201  
 3204  
 3207  
 3210  
 3213  
 3216  
 3219  
 3222  
 3225  
 3228  
 3231  
 3234  
 3237  
 3240  
 3243  
 3246  
 3249  
 3252  
 3255  
 3258  
 3261  
 3264  
 3267  
 3270  
 3273  
 3276  
 3279  
 3282  
 3285  
 3288  
 3291  
 3294  
 3297  
 3300  
 3303  
 3306  
 3309  
 3312  
 3315  
 3318  
 3321  
 3324  
 3327  
 3330  
 3333  
 3336  
 3339  
 3342  
 3345  
 3348  
 3351  
 3354  
 3357  
 3360  
 3363  
 3366  
 3369  
 3372  
 3375  
 3378  
 3381  
 3384  
 3387  
 3390  
 3393  
 3396  
 3399  
 3402  
 3405  
 3408  
 3411  
 3414  
 3417  
 3420  
 3423  
 3426  
 3429  
 3432  
 3435  
 3438  
 3441  
 3444  
 3447  
 3450  
 3453  
 3456  
 3459  
 3462  
 3465  
 3468  
 3471  
 3474  
 3477  
 3480  
 3483  
 3486  
 3489  
 3492  
 3495  
 3498  
 3501  
 3504  
 3507  
 3510  
 3513  
 3516  
 3519  
 3522  
 3525  
 3528  
 3531  
 3534  
 3537  
 3540  
 3543  
 3546  
 3549  
 3552  
 3555  
 3558  
 3561  
 3564  
 3567  
 3570  
 3573  
 3576  
 3579  
 3582  
 3585  
 3588  
 3591  
 3594  
 3597  
 3600  
 3603  
 3606  
 3609  
 3612  
 3615  
 3618  
 3621  
 3624  
 3627  
 3630  
 3633  
 3636  
 3639  
 3642  
 3645  
 3648  
 3651  
 3654  
 3657  
 3660  
 3663  
 3666  
 3669  
 3672  
 3675  
 3678  
 3681  
 3684  
 3687  
 3690  
 3693  
 3696  
 3699  
 3702  
 3705  
 3708  
 3711  
 3714  
 3717  
 3720  
 3723  
 3726  
 3729  
 3732  
 3735  
 3738  
 3741  
 3744  
 3747  
 3750  
 3753  
 3756  
 3759  
 3762  
 3765  
 3768  
 3771  
 3774  
 3777  
 3780  
 3783  
 3786  
 3789  
 3792  
 3795  
 3798  
 3801  
 3804  
 3807  
 3810  
 3813  
 3816  
 3819  
 3822  
 3825  
 3828  
 3831  
 3834  
 3837  
 3840  
 3843  
 3846  
 3849  
 3852  
 3855  
 3858  
 3861  
 3864  
 3867  
 3870  
 3873  
 3876  
 3879  
 3882  
 3885  
 3888  
 3891  
 3894  
 3897  
 3900  
 3903  
 3906  
 3909  
 3912  
 3915  
 3918  
 3921  
 3924  
 3927  
 3930  
 3933  
 3936  
 3939  
 3942  
 3945  
 3948  
 3951  
 3954  
 3957  
 3960  
 3963  
 3966  
 3969  
 3972  
 3975  
 3978  
 3981  
 3984  
 3987  
 3990  
 3993  
 3996  
 3999  
 4002  
 4005  
 4008  
 4011  
 4014  
 4017  
 4020  
 4023  
 4026  
 4029  
 4032  
 4035  
 4038  
 4041  
 4044  
 4047  
 4050  
 4053  
 4056  
 4059  
 4062  
 4065  
 4068  
 4071  
 4074  
 4077  
 4080  
 4083  
 4086  
 4089  
 4092  
 4095  
 4098  
 4101  
 4104  
 4107  
 4110  
 4113  
 4116  
 4119  
 4122  
 4125  
 4128  
 4131  
 4134  
 4137  
 4140  
 4143  
 4146  
 4149  
 4152  
 4155  
 4158  
 4161  
 4164  
 4167  
 4170  
 4173  
 4176  
 4179  
 4182  
 4185  
 4188  
 4191  
 4194  
 4197  
 4200  
 4203  
 4206  
 4209  
 4212  
 4215  
 4218  
 4221  
 4224  
 4227  
 4230  
 4233  
 4236  
 4239  
 4242  
 4245  
 4248  
 4251  
 4254  
 4257  
 4260  
 4263

C TAG AAA AAA CCA AGG AGG TAA TAA ATA 30 ATG GCA CCT TTA GGT CCA ACT GGT CCT CTG 60  
 Met Ala Pro Leu Gly Pro Thr Gly Pro Leu  
 CCT CAA AGT TTC CTG CTG AAA TGC CTC GAG CAG ATG CCG AAA GTT CAA GCT GAT GGT ACC 120  
 Pr Gln Ser Phe Leu Leu Lys Cys Leu Glu Gln Met Arg Lys Val Gln Ala Asp Gly Thr  
 GCA CTC CAA GAA ACT CTG TGC GCA ACT CAC CAA CTG TGC CAC CCT GAA GAA CTC GTA CTG 180  
 Ala Leu Gln Glu Thr Leu Cys Ala Thr His Gln Leu Cys His Pro Glu Glu Leu Val Leu  
 CTC GGT CAC GCA CTC GGT ATT CCG CAG CCG CTC TCT TCT TCT TGC TCC TCT CAG GCT CTG 240  
 Leu Gly His Ala Leu Gly Ile Pro Gln Pro Pro Leu Ser Ser Ser Cys Ser Ser Gln Ala Leu  
 CAA CTC ATG GGT TGC CTC CGT CAA CTG CAT TCT GGC CTG TTC CTG TAC CAG GGT CTC CTG 300  
 Gln Leu Met Gly Cys Leu Arg Gln Leu His Ser Gly Leu Phe Leu Tyr Gln Gly Leu Leu

FIG. 4-A

8 / 16

330 CAA GCT TTG GCT GGC ATC TCT CCG GAA CTC GCA CCT ACT CTC GAC ACT CTG CAG CTC GAC 360  
 Gln Ala Leu Ala Gly Ile Ser Pro Glu Leu Ala Pro Thr Leu Asp Thr Leu Gln Leu Asp  
 390 ACT ACC GAC TTC GCT ATC AAC ATT TGG CAG CAA ATG GAA GAT CTG GGC ATG GCA CCG GCT 420  
 Thr Thr Asp Phe Ala Ile Asn Ile Trp Gln Gln Met Glu Asp Leu Gly Met Ala Pro Ala  
 450 GTT CCG CCG ACT CAG GGC ACT ATG CCT GCT TTT ACT TCT GCT TTC CAG CGT CGT GCT GGT 480  
 Val Pro Pro Thr Gln Gly Thr Met Pro Ala Phe Thr Ser Ala Phe Gln Arg Arg Ala Gly  
 510 GGT GTA CTC GTA GCT TCT AAC CTC CAG TCT TTC CTC GAA CTC GCT TAC CGT GCT CTG CGT 540  
 Gly Val Leu Val Ala Ser Asn Leu Gln Ser Phe Leu Glu Leu Ala Tyr Arg Ala Leu Arg  
 CAC TTC GCT AAA CCG TAA TAG GAT C  
 His Phe Ala Lys Pro End End Asp

FIG. 4-B

SUBSTITUTE SHEET



9 / 16

```

10  CTAGAAA AACCAAGGAG GTAATAATA ATGGCACCTT TAGGTCCAAC TGGTCCTCTG 60
    xba1 TTTT TTGGTTCCCTC CATTATTAT TACCGTGGAA ATCCAGGTTG ACCAGGAGAC
70  CCTCAAAGTT TCCTGCTGAA ATGCCCTCGAG CAGATGCGTA AAGTTC AAGC TGA TGGTACC 120
    GGAGTTTCAA AGGACGACTT TACGGAGCTC GTCTACGCAT TTCAAGTTTCG ACTACCATGG
130  GCACTCCAAG AACTCTGTG CGCAACTCAC CAACTGTGCC ACCCTGAAGA ACTCGTACTG 180
    CGTGAGGTTT TTTGAGACAC GCGTTGAGTG GTTACACCGG TGGGACTTCT TGAGCATGAC
190  CTCGGTCACG CACTCGGTAT TCCGCAGCCG CCGCTGTCTT CTGCTCCTC TCAGGCTCTG 240
    GAGCCAGTGC GTGAGCCATA AGGCGTCGGC GCGACAGAA GAACGAGGAG AGTCCGAGAC
250  CAACTCATGG GTTGCCTCCG TCAACTGCAT TCTGGCCCTGT TCCTGTACCA GGTCTCCTG 300
    GTTGAGTACC CAACGGAGGC AGTTGACGTA AGACCGGACA AGGACATGGT CCCAGAGGAC

```

FIG. 5-A

SUBSTITUTE SHEET

10 / 16

```

310 320 330 340 350 360
CAAGCTTTGG CTGGCATCTC TCCGGAAC TC CACTTACTC TCGACACTCT GCAGCTCGAC
GTTCCGAAACC GACCGTAGAG AGGCCTTGAG CGTGGATGAG AGCTGTGAGA CGTCGAGCTG
HindIII
370 380 390 400 410 420
ACTACCGACT TCGCTATCAA CATTTGGCAG CAAATGGAAG ATCTGGGCAT GGCACCGGCT
TGATGGCTGA AGCGATAGTT GTAAACCGTC GTTTACCTTC TAGACCCGTA CCGTGGCCGA
430 440 450 460 470 480
GTTCCGCCGA CTCAGGGCAC TATGCCCTGCT TTTACTTCTG CTATCCAGCG TCGTGCTGGT
CAAGGCGGCT GAGTCCGTG ATACGGACGA AAATGAAGAC GAAAGGTCGC AGCACGACCA
490 500 510 520 530 540
GGTGACTCG TAGCTTCTAA CCTCCAGTCT TTCCCTCGAAC TCGCTTACCG TGCTCTGCCG
CCACATGAGC ATCGAAGATG GGAGGTCAGA AAGGAGCTTG AGCGAATGGC ACAGAGACGCA
550 560
CACTTCGCTA AACCGTAATA G BamHI
GTGAAGCGAT TTGGCATTAT CCTAG

```

FIG. 5-B

SUBSTITUTE SHEET

11 / 16

```

10      20      30      40      50      60
CTAGAAAAA CCAAGGAGGT AATAAATAAT GGCACCTTAA GGTCCAAC TG GTCTCTGCCC
xbal TTTTTT GGTTCCTCCA TTATTATTA CCGTGGAAAT CQAGGTTGAC CAGGAGACGG

70      80      90      100     110     120
TCAAAGTTTC CTGCTGAAAT GCCTCGAGCA GATGCGTAAA GTTCAAGCTG ATGGTACCGC
AGTTTCAAAG GACGACTTTA CGGAGCTCGT CTACGCATTT CAAGTTTCGAC TACCATGGCG

130     140     150     160     170     180
ACTCCAAGAA ACTCTGTGCG CAACTCAACA ACTGTGCCAC CCTGAAGAAC TCGTACTGCT
TGAGGTTCTT TGAGACACGC GTTGAGTGGT TGACACGGTG GGACTTCTTG AGCATGACGA

190     200     210     220     230     240
CGGTTCACGCA CTCGGTATTG CGCAGCCGCC GCTGTCTTCT TGCTCTCTC AGGCTCTGCA
GCCAGTGCGT GAGCCATAAG GCGTCGGCGG CGACAGAAGA ACGAGGAGAG TCCGAGACGT

250     260     270     280     290     300
ACTCATGGGT TGCCTCGTC AACTGCATTG TGGCCTGTTC CTGTACCAGG GTCTCCGTCA HindIII
TGAGTACCCA ACGGAGGCAG TTGACGTAAG ACCGGACAAG GACATGGTCC CAGAGGACGT TCGA

```

FIG. 6-A

12 / 16

```

10  AGCTTTGGCT GGCATCTCTC CGGAACCTCGC ACCTACTCTC GACACTCTGC AGCTCGACAC 60
    AACCGA CCGTAGAGAG GCCTTGAGCG TGGATGAGAG CTGTGAGACG TCGAGCTGTG
    HindIII
70  TACCCACTTC GCTATCAACA TTTGGCAGCA AATGAAGAT CTGGGCATGG CACCGGCTGT 120
    ATGGCTGAG CGATAGTTGT AAACCGTCGT TTACCTTCTA GACCCGTACC GTGGCCGACA
130  TCCGCCGACT CAGGGCACTA TGCCTGCTTT TACTTCTGCT TTCCAGCGTC GTGCTGGTGG 180
    AGCGGGCTGA GTCCTCGTAT ACGGACGAAA ATGAAGACGA AAGGTCGCAG CACGACCACC
190  TGTAATCGTA GCTTCTAACC TCCAGTCTTT CCTCGAATC GCTTACCGTG CTCTGCGTCA 240
    ACATGAGCAT CGAAGATTGG AGGTCAGAAA GGAGCTTGAG CGAATGGCAC GAGACGCAGT
250  CTTTCGCTAA CCGTAATAG BamHI
    GAAGCGATTT GGCAATTATCC TAG
260

```

FIG. 6-B

13 / 16

```

1      MetAlaProLeuGlyProThrGlyProLeuProGlnSerPheLeuLeuLysCysLeuGlu
      | : | | | | : | | | | | | | | | | |
1      MetThrProLeuGlyProAlaSerSerLeuProGlnSerPheLeuLeuLysCysLeuGlu

21     GlnMetArgLysValGlnAlaAspGlyThrAlaLeuGlnGluThrLeuCysAlaThrHis
      | : | | | | : | | | | | | | | | | |
21     GlnValArgLysIleGlnGlyAspGlyAlaAlaGluGlnGluLysLeuCysAlaThrTyr

41     GlnLeuCysHisProGluGluLeuValLeuLeuGlyHisAlaLeuGlyIleProGlnPro
      | | | | | | | | | | | | : | | | | |
41     LysLeuCysHisProGluGluLeuValLeuLeuGlyHisSerLeuGlyIleProTrpAla

61     ProLeuSerSerCysSerSerGlnAlaLeuGlnLeuMetGlyCysLeuArgGlnLeuHis
      | | | | | | | | | | | | | | | | |
61     ProLeuSerSerCysProSerGlnAlaLeuGlnLeuAlaGlyCysLeuSerGlnLeuHis

81     SerGlyLeuPheLeuTyrGlnGlyLeuLeuGlnAlaLeuAlaGlyIleSerProGluLeu
      | | | | | | | | | | | | | | | | |
81     SerGlyLeuPheLeuTyrGlnGlyLeuLeuGlnAlaLeuGluGlyIleSerProGluLeu

```

FIG. 7-A

14 / 16

```

101 AlaProThrLeuAspThrLeuGlnLeuAspThrThrAspPheAlaIleAsnIleTrpGln
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
101 GlyProThrLeuAspThrLeuGlnLeuAspValAlaAspPheAlaThrThrIleTrpGln
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
121 GlnMetGluAspLeuGlyMetAlaProAlaValProProThrGlnGlyThrMetProAla
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
121 GlnMetGluGluLeuGlyMetAlaProAlaLeuGlnProThrGlnGlyAlaMetProAla
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
141 PheThrSerAlaPheGlnArgArgAlaGlyGlyValLeuValAlaSerAsnLeuGlnSer
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
141 PheAlaSerAlaPheGlnArgArgAlaGlyGlyValLeuValAlaSerHisLeuGlnSer
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
161 PheLeuGluLeuAlaTyrArgAlaLeuArgHisPheAlaLysPro
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |
161 PheLeuGluValSerTyrArgValLeuArgHisLeuAlaGlnPro
      | | | | | | | | | | | | | | | | | | | | | | | | | | | |

```

FIG. 7-B

# **Canine G-CSF Study** Neutrophil Counts




FIG. 8

**Canine G-CSF Study**  
Mean Counts (n=2)




FIG. 9



# INTERNATIONAL SEARCH REPORT

International Application No PCT/US90/05522

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>3</sup> According to International Patent Classification (IPC) or to both National Classification and IPC IPC(5): C07K 13/00; A61K 37/02; C12P 21/02; C12N 15/24; C07H 15/12 U.S.CL.: 530/351; 435/69.5,172.3,240.2,243,320; 536/27; 424/85.1																				
<b>II. FIELDS SEARCHED</b> <div style="text-align: center; font-size: small;">Minimum Documentation Searched <sup>4</sup></div> <table style="width: 100%; border: none;"> <tr> <td style="width: 30%; border: none;">Classification System</td> <td style="border: none;">Classification Symbols</td> </tr> <tr> <td style="border: none;">U.S.CL.</td> <td style="border: none;">530/351,395,300,820,827; 435/69.5,172.3,240.2,243,320; 424/85.1; 514/2,8,12; 536/27</td> </tr> </table>			Classification System	Classification Symbols	U.S.CL.	530/351,395,300,820,827; 435/69.5,172.3,240.2,243,320; 424/85.1; 514/2,8,12; 536/27														
Classification System	Classification Symbols																			
U.S.CL.	530/351,395,300,820,827; 435/69.5,172.3,240.2,243,320; 424/85.1; 514/2,8,12; 536/27																			
<div style="text-align: center; font-size: x-small;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>5</sup></div>																				
<b>COMPUTER DATA-BASE SEARCH ON CAS FOR: CANINE OR FELINE AND G-CSF, AND (DNA OR RECOMBINANT)</b>																				
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>14</sup></b> <table border="1" style="width: 100%; border-collapse: collapse; font-size: small;"> <thead> <tr> <th style="width: 10%;">Category <sup>6</sup></th> <th style="width: 70%;">Citation of Document, <sup>15</sup> with indication, where appropriate, of the relevant passages <sup>17</sup></th> <th style="width: 20%;">Relevant to Claim No. <sup>18</sup></th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">Y</td> <td>US, A, 4,833,127 (Ono et al.) 23 May 1989, see entire document.</td> <td style="text-align: center;">1-39</td> </tr> <tr> <td style="text-align: center;">Y</td> <td>US, A, 4,810,643 (Souza) 07 March 1989, see entire document.</td> <td style="text-align: center;">1-39</td> </tr> <tr> <td style="text-align: center;">Y, P</td> <td>US, A, 4,904,584 (Shaw) 27 February 1990 see entire document.</td> <td style="text-align: center;">1-39</td> </tr> <tr> <td style="text-align: center;">Y</td> <td>EP, A, 0,220,520 (Yamozaki) 05 June 1987, see the entire document.</td> <td style="text-align: center;">1-39</td> </tr> <tr> <td style="text-align: center;">Y</td> <td>Proc. Natl. Acad. Sci., Vol. 83, Issued October 1986, Tsuchiya et al., "Isolation and Characterization of the cDNA for Murine Granulocyte Colony-Stimulating Factor", pages 7633-37, (see pages 7633 and 7635-30).</td> <td style="text-align: center;">1-39</td> </tr> </tbody> </table>			Category <sup>6</sup>	Citation of Document, <sup>15</sup> with indication, where appropriate, of the relevant passages <sup>17</sup>	Relevant to Claim No. <sup>18</sup>	Y	US, A, 4,833,127 (Ono et al.) 23 May 1989, see entire document.	1-39	Y	US, A, 4,810,643 (Souza) 07 March 1989, see entire document.	1-39	Y, P	US, A, 4,904,584 (Shaw) 27 February 1990 see entire document.	1-39	Y	EP, A, 0,220,520 (Yamozaki) 05 June 1987, see the entire document.	1-39	Y	Proc. Natl. Acad. Sci., Vol. 83, Issued October 1986, Tsuchiya et al., "Isolation and Characterization of the cDNA for Murine Granulocyte Colony-Stimulating Factor", pages 7633-37, (see pages 7633 and 7635-30).	1-39
Category <sup>6</sup>	Citation of Document, <sup>15</sup> with indication, where appropriate, of the relevant passages <sup>17</sup>	Relevant to Claim No. <sup>18</sup>																		
Y	US, A, 4,833,127 (Ono et al.) 23 May 1989, see entire document.	1-39																		
Y	US, A, 4,810,643 (Souza) 07 March 1989, see entire document.	1-39																		
Y, P	US, A, 4,904,584 (Shaw) 27 February 1990 see entire document.	1-39																		
Y	EP, A, 0,220,520 (Yamozaki) 05 June 1987, see the entire document.	1-39																		
Y	Proc. Natl. Acad. Sci., Vol. 83, Issued October 1986, Tsuchiya et al., "Isolation and Characterization of the cDNA for Murine Granulocyte Colony-Stimulating Factor", pages 7633-37, (see pages 7633 and 7635-30).	1-39																		
<div style="display: flex; justify-content: space-between; font-size: x-small;"> <div style="width: 45%;"> <p><b>* Special categories of cited documents: <sup>15</sup></b></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&amp;" document member of the same patent family</p> </div> </div>																				
<b>IV. CERTIFICATION</b> <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none;">           Date of the Actual Completion of the International Search <sup>1</sup>   <div style="text-align: center; font-size: large;">13 December 1990</div> </td> <td style="width: 50%; border: none;">           Date of Mailing of this International Search Report <sup>2</sup>   <div style="text-align: center; font-size: x-large;">04 FEB 1991</div> </td> </tr> <tr> <td style="border: none;">           International Searching Authority <sup>1</sup>   <div style="text-align: center; font-size: large;">ISA/US</div> </td> <td style="border: none;">           Signature of Authorized Officer <sup>19</sup>    <div style="text-align: center; font-weight: bold;">GARNETTE D. DRAPER, PRIMARY EXAMINER</div> </td> </tr> </table>			Date of the Actual Completion of the International Search <sup>1</sup>  <div style="text-align: center; font-size: large;">13 December 1990</div>	Date of Mailing of this International Search Report <sup>2</sup>  <div style="text-align: center; font-size: x-large;">04 FEB 1991</div>	International Searching Authority <sup>1</sup>  <div style="text-align: center; font-size: large;">ISA/US</div>	Signature of Authorized Officer <sup>19</sup>  <div style="text-align: center; font-weight: bold;">GARNETTE D. DRAPER, PRIMARY EXAMINER</div>														
Date of the Actual Completion of the International Search <sup>1</sup>  <div style="text-align: center; font-size: large;">13 December 1990</div>	Date of Mailing of this International Search Report <sup>2</sup>  <div style="text-align: center; font-size: x-large;">04 FEB 1991</div>																			
International Searching Authority <sup>1</sup>  <div style="text-align: center; font-size: large;">ISA/US</div>	Signature of Authorized Officer <sup>19</sup>  <div style="text-align: center; font-weight: bold;">GARNETTE D. DRAPER, PRIMARY EXAMINER</div>																			

## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Y

The EMBO Journal, Vol. 6, No. 3, Issued 1987, (Tsuchiya et al.), "Characterization of Recombinant Human Granulocyte-Colony-Stimulating Factor Produced in Mouse Cells", pages 611-616, (see pages 611 and 614-15).

1-39

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE<sup>1</sup>

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers ..... because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claim numbers ..... because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out<sup>1</sup>, specifically:

3. ☐ Claim numbers ..... because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING<sup>2</sup>

This international Searching Authority found multiple inventions in this international application as follows: Group I, Claims 1-8, 11-12, 25-27, 39 to polypeptide to G-CSF and composition; classified in 530/351. Group II, claims 9-10, 13-24 and 28 to DNA, cell lines plasmid and recombinant production of protein, classified in 435/69.5, 172.3. Group III, claims 29-38 to a method of treatment, classified in 424/85.1.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☒ As all searchable claims could be searched without effort justifying an additional fee, the international Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.  
☐ No protest accompanied the payment of additional search fees.